ple, 2.04 of resin, 1.38 of tannin, a slight portion of gum, and the rest of fibrina and earthy salts. The resin is adhesive, brownish, readily soluble in ether and alkalis, burning with flame and a resinous odour, and leaving a white cinder.

From my own trials the quantity of resin in this plant appears to be very small. A saturated tincture of a deep brown colour does not give a precipitate on the first addition of water. It is only after some time standing, and partly perhaps from the evaporation of the alcohol, that a turbidness begins to appear in the solution. It is probable that spirit is a better menstruum than water for the soluble portions of this plant, although the latter is capable of extracting the greater part of its virtue.

The Pyrola umbellata, though scarcely known as a medicine until within a few years past, has at the present day acquired a reputation of considerable extent in the treatment of various diseases. Its popular celebrity seems to have originated in its application to the treatment of fever and rheumatism; but the attention of physicians has been chiefly drawn towards its use in other complaints. The instances in which this plant has received favourable testimonies on medical authority, of its successful use, both in America

and Europe, are principally the following. 1. As a palliative in strangury and nephritis. 2. As a diuretic in dropsy. 3. As an external stimulant, susceptible of useful application to various cases.

In the first of these cases, the Pyrola is entitled to attention and confidence. Some practitioners in this country have employed it with advantage in the same cases, in which the Arbutus Uva ursi is recommended*. Dr. Wolf, the German writer lately cited, has reported a number of cases of ischuria and dysuria, arising from various causes, in which the Pyrola, given in infusion, produced the most evident relief, and took precedence of a variety of remedies which had been tried. His method of administering it was to give a table spoonful of a strong infusion, with a little syrup, every hour. In all the cases he has detailed, small as the dose was, it gave relief in a very short time. In one case its effect was so distinctly marked, that the disease returned whenever the medicine was omitted and was removed on resuming its use. A tonic operation attended its other effects, so that the appetite was improved and digestion promoted during the period of its employment.

* See Dr. Mitchell's Inaugural Dissertation. Philadelphia, 1803.